



ANTHRAX e-TOOL Protecting the Worksite against terrorism

| | | | | | |
|----------------------|----------------------------------|---------------------------------|-----------------------------------|--|--|
| Home | What is anthrax? | Who is at risk? | How do I prepare? | What if I have a threat? | How do I clean up my worksite? |
|----------------------|----------------------------------|---------------------------------|-----------------------------------|--|--|

How do I clean up my worksite that has been contaminated with anthrax?

The following pages were developed as a technical resource specifically for response to a release of anthrax. The purpose of these pages is to help protect public health and safety by providing the most current information available throughout the federal government, and sharing national experience to date in responding to releases of *Bacillus anthracis* in urban environments.



- ④ [What first response actions should be taken?](#) This section describes activities during the initial or emergency phase (generally the first 24-48 hours) of a response to a suspected incident involving anthrax. It is intended for personnel who discover the potential contamination and for first responders on the scene.
- ④ [What Health and Safety Plan \(HASP\) requirements exist at anthrax contaminated sites?](#) This section describes the Health and Safety Plan requirements as they apply to response activities related to an anthrax release.
- ④ [What training requirements exist for workers involved in anthrax response and remediation?](#) This section describes the training that is required to provide response workers and other potentially affected persons the hazard awareness training they need to work safely.
- ④ [What types of personal protective equipment \(PPE\) are necessary?](#) This section describes the Personal Protective Equipment (PPE) necessary to shield or isolate workers from health and safety hazards in the workplace. In a site where anthrax spores may be present, PPE protects workers from exposure to respiratory and skin hazards and prevents the spread of contaminants to uncontaminated areas.
- ④ [What are the employee medical program requirements?](#) This section describes the Medical Surveillance Program requirements for response activities related to an anthrax release.
- ④ [How do I sample and analyze for anthrax?](#) This section describes sampling methodologies for anthrax releases and analytical methods for detection of anthrax.
- ④ [How should I decontaminate during response actions?](#) This section provides information on decontaminating buildings or specific areas, systems, or items within buildings after an actual release of anthrax.
- ④ [What is a Transition Program and what elements should I include in my Transition Program?](#) This section describes the Transitional Program requirements for facilities in which anthrax contamination has been identified, response actions have been completed, and normal work operations will be conducted.
- ④ [Links to Additional Anthrax Response Information](#) This web page contains links to references and additional information for each of the sections above.



ANTHRAX e-TOOL Protecting the Worksite against terrorism

- Home
- What is anthrax?
- Who is at risk?
- How do I prepare?
- What if I have a threat?
- How do I clean up my worksite?

What first response actions should be taken?

This page describes activities during the initial or emergency phase (generally the first 24-48 hours) of a response to a suspected incident involving anthrax. It is intended for personnel who discover the potential contamination and for first responders on the scene.



- ④ [Notification and Initial Actions](#)
- ④ [Lead Agencies and Incident Command System/Unified Command](#)
- ④ [Preliminary Assessment to Determine Credible Threat](#)
- ④ [Site Assessment to Verify Contamination](#)
- ④ [Identification of Potentially Exposed Personnel](#)

Notification and Initial Actions

As soon as a initial anthrax contamination is confirmed:

- ④ Call **9-1-1** for local police assistance and the local Federal Bureau of Investigation (FBI). Federal agencies located in buildings managed by GSA should contact the Federal Protective Service.
- ④ Notify the **National Response Center (NRC) at (800)424-8802**. The NRC will notify appropriate agencies, such as the Environmental Protection Agency (EPA), FBI, United States Coast Guard (USCG), and Centers for Disease Control and Prevention (CDC), to arrange needed assistance.
- ④ Contact the owner or operator of the facility.
- ④ Direct occupants of the facility to report to a designated meeting place, as identified in the local facility response plan.



Immediately dial 9-1-1 if you have a confirmed anthrax attack.

Lead Agencies and Incident Command System/Unified Command

The lead agencies responding to a suspected incident will coordinate their efforts under a Unified Command (UC) within an Incident Command System (ICS) to ensure their actions and resources are integrated into a single, effective response. The UC for anthrax response will usually include:

- ④ FBI
- ④ Federal On-scene Coordinator (such as the EPA, USCG)
- ④ State On-scene Coordinator and/or local Incident

Commander(s) (such as the local police, fire department, or public health official)

- ④ Facility manager or property owner



Preliminary Assessment to Determine Credible Threat

Initial efforts should be made to assess the situation and determine whether the incident poses a credible threat. The assessment may begin with a conference call with the reporting party and involved agencies, such as the FBI, EPA, USCG, CDC, and appropriate local health and law enforcement authorities. If the report appears to be credible, the FBI and other responding agencies will coordinate a site assessment to confirm the threat.



Site Assessment to Verify Contamination

The objectives of the Site Assessment sampling may include any of the following:

- ④ **Determine qualitatively whether any spores are present.** Typically, composite samples of large areas and air volumes are obtained to maximize the likelihood of finding contamination.
- ④ **Identification of Spores in a Bulk Material (such as powder in an envelope)** On-site analysis may be used for preliminary assessment to determine qualitatively if a bulk material is contaminated with anthrax.
- ④ **Determination of Contamination of an Article** Composite surface samples of large articles and individual samples of small articles are collected to determine whether the article's surface is contaminated.



Vacuums such as this are used to sample large volumes of air.

Identification of Potentially Exposed Personnel

As quickly as possible, facility managers and first responders should identify personnel who might have been exposed, including:

- ④ People in the area where the spores were released,
- ④ People who may have handled contaminated items or packages, and
- ④ Response personnel.

Potential sources of data include:

- Ⓢ Employee timecards,
- Ⓢ Visitor logs, and
- Ⓢ Security videotapes.

| [Home](#) | [What is Anthrax?](#) | [Who is at risk?](#) | [How do I prepare?](#) |
| [What if I have an anthrax threat?](#) | [How do I clean up my worksite?](#) |
| [Viewing/Printing Instructions](#) | [Credits](#) |

 [Back to Top](#)

www.osha.gov

www.dol.gov

[Contact Us](#) | [Freedom of Information Act](#) | [Information Quality](#) | [Customer Survey](#)
[Privacy and Security Statement](#) | [Disclaimers](#)

Occupational Safety & Health Administration
200 Constitution Avenue, NW
Washington, DC 20210



ANTHRAX eTool.....Protecting the Worksite against terrorism

| | | | | | |
|----------------------|----------------------------------|---------------------------------|-----------------------------------|--|--|
| Home | What is anthrax? | Who is at risk? | How do I prepare? | What if I have a threat? | How do I clean up my worksite? |
|----------------------|----------------------------------|---------------------------------|-----------------------------------|--|--|

What Health and Safety Plan (HASP) requirements exist for anthrax-contaminated sites?

For cleanup operations, OSHA's *Hazardous Waste Operations and Emergency Response (HAZWOPER)* standard (29 CFR 1910.120) requires a written health and safety plan (HASP), which identifies site hazards and appropriate controls to protect employee health and safety. The elements of the HASP are described in the standard and include the following:



- ④ [Organizational Structure](#)
- ④ [Site Characterization and Job Hazard Analysis](#)
- ④ [Site Control](#)
- ④ [Training](#)
- ④ [Medical Surveillance](#)
- ④ [Personal Protective Equipment](#)
- ④ [Exposure Monitoring](#)
- ④ [Temperature Extremes](#)
- ④ [Spill Containment](#)
- ④ [Decontamination](#)
- ④ [Emergency Response](#)
- ④ [Standard Operating Procedures](#)

Due to overlap of some of the elements, it may be useful to expand the HASP to include those elements necessary to protect the local community and environment (for example, disposal of waste from decontamination, monitoring community exposures to fumigants).

Organizational Structure

The organizational structure part of the program shall establish the specific chain of command and specify the overall responsibilities of supervisors and employees. It shall include, at a minimum, the following elements:

- ④ A general supervisor who has the responsibility and authority to direct all hazardous waste operations.
- ④ A site safety and health supervisor who has the responsibility and authority to develop and implement the HASP and verify compliance.
- ④ All other personnel needed for hazardous waste site operations and emergency response and their general functions and responsibilities.
- ④ The lines of authority, responsibility, and communication.

The organizational structure shall be reviewed and updated as necessary to reflect the current status of waste site operations. It is necessary to organize and assign personnel roles, responsibilities, lines of authority, and communication before an anthrax-related event to ensure an efficient and safe response.

For more detailed information regarding specific roles associated with an emergency response, go to the [Anthrax eTool Training page](#).

Site Characterization and Job Hazard Analysis

The HASP requires a characterization of the site-specific job hazards, including:

- ④ Location and approximate size of the site,
- ④ Description of the response activity and the job task to be performed,
- ④ Duration of the planned employee activity,
- ④ Site topography and accessibility by air and roads,
- ④ Safety and health hazards expected at the site (for example, chemical, physical, biological hazards),
- ④ Pathways for hazardous substance spread, and
- ④ Present status and capabilities of emergency response teams that would provide assistance to hazardous waste cleanup site employees at the time of an emergency.



The following links provide additional information regarding site characterization and job hazard analysis:

- ④ [Anthrax eTool: Who is at risk of anthrax exposure?](#)
- ④ [OSHA's Hazardous Waste Operations and Emergency Response \(HAZWOPER\) Standard 29 CFR 1910.120](#)

Site Control

In the event of an anthrax release, it is necessary to control site access in order to prevent exposure and spread of the anthrax spores. Suggested site controls include:

- ④ Evacuate and secure the area,
- ④ Turn off fans and air handling systems, if possible, and
- ④ Notify personnel about the potential hazard and to not access the area. The impacted area should only be accessed by those trained and informed about the release.



As outlined in the [HAZWOPER \(29 CFR 1910.120\)](#), a site control program for protecting employees should be part of the employer's HASP and should include, at a minimum, the following:

- ④ Site map,
- ④ Site work zones,
- ④ Use of a "buddy system,"
- ④ Site communications including alerting means for emergencies,
- ④ Standard Operating Procedures (SOPs) or safe work practices, and

- ④ Identification of the nearest medical assistance.

Training

A site-specific training program ensures that workers receive the hazard awareness training they need to work safely. Training should be based on the job hazard analysis in the HASP and other applicable standards.

Anthrax-specific hazard awareness training should help workers understand the health hazards of anthrax and how to protect themselves from exposure to spores. Specific topics might include:

- ④ How workers might be exposed to spores, the signs and [symptoms of infection](#), and medical conditions that could place them at increased risk (such as a compromised immune system),
- ④ Where contamination has been identified in the facility, and the status of [decontamination](#) of those areas, and
- ④ How to minimize the risk of disease through specific standard operating procedures and controls (such as engineering controls, work practices, housekeeping, or [PPE](#)), and whether specific measures are expected to be temporary or permanent.



49 CFR Part 172, Subpart H provides additional training requirements for workers preparing contaminated materials or other hazardous materials for transport. For the most current version of 49 CFR, visit the: [Code of Federal Regulations website](#).

Additional training information is provided on the [Anthrax eTool Training page](#).

Medical Surveillance

The purpose of medical surveillance in the workplace is to improve the effectiveness of the occupational health and safety program by systematically collecting and analyzing information that pertains to at-risk workers.

Medical screening is the use of examinations or tests to detect adverse effects on a worker's health at an early stage when prevention is possible or treatment is most effective.

- ④ **Baseline medical screening** should identify pre-existing conditions that may affect an individual worker's fitness for duty.
- ④ When it is no longer necessary for a worker to re-enter a contaminated site, a **final evaluation** should be done to identify changes from the baseline and any new risk factors.



Further information about Medical Surveillance may be found at the following websites:

- ④ Centers for Disease Control and Prevention (CDC), Morbidity and Mortality Weekly Report (MMWR), [Notice to Readers: Occupational Health Guidelines for Remediation Workers at Bacillus anthracis-Contaminated Sites \[September 6, 2002 / 51\(35\);786-789\]](#)

Personal Protective Equipment

[Personal protective equipment](#) (PPE) shields or isolates workers from health and safety hazards in the workplace. In a site where anthrax spores may be present, PPE protects workers from exposure to respiratory and skin hazards and prevents the spread of contaminants to uncontaminated areas. The level and type of PPE should be based on the job hazard analysis in the HASP. Workers should be trained on the location, type, and proper use of the PPE.



Further information about PPE may be found at the following websites:

- ④ [OSHA's PPE Safety and Health Topics page](#)

Exposure Monitoring

For exposure monitoring, the HASP should include the following three major components:

- ④ Medical measures to prevent anthrax,
- ④ Medical screening and follow-up care for anthrax and medical complications related to preventive measures, and
- ④ Knowledge and information that workers need to prevent anthrax and medical complications related to preventive measures.

Temperature Extremes

Heat Stress: Temperature extremes can adversely affect worker health and safety. Heat stress can be experienced by workers due to hot ambient temperatures. In addition, the use of PPE, especially in excess, can increase the chance a worker will experience heat stress. The four main types of heat stress are listed below:

- ④ **Heat Rash:** Caused by continuous exposure to heat and humid air and aggravated by chafing clothing. Decreases ability to tolerate heat, as well as being a nuisance.
- ④ **Heat Cramps:** Caused by profuse perspiration with inadequate fluid intake and chemical replacement. Signs are muscle spasms and pain in the extremities and abdomen.
- ④ **Heat Exhaustion:** Caused by increased stress on various organs to meet increased demands to cool the body. Signs are shallow breathing; pale, cool, moist skin; profuse sweating; and dizziness and physical or mental exhaustion.
- ④ **Heat Stroke:** The most severe form of heat stress. Body must be cooled immediately to prevent severe injury and/or death. Signs are red, hot, dry skin; no perspiration; nausea; dizziness and confusion; strong, rapid pulse; coma. Medical help must be obtained immediately.



Cold Stress: Cold stress can be experienced by workers when they are exposed to a cold environment. The two main types of cold stress are listed below:

- ④ **Frostbite:** Local injury resulting from cold is included in the generic term frostbite. There are

several degrees of damage. Frostbite of the extremities can be categorized into:

1. *Frost nip or incident frostbite* - the condition is characterized by sudden blanching or whitening of skin.
2. *Superficial frostbite* - skin has a waxy or white appearance and is firm to the touch, but tissue beneath is resilient.
3. *Deep frostbite* - tissues are cold, pale and solid; extremely serious injury.

④ **Hypothermia:** There are degrees of hypothermia which are characterized as "moderate" and "severe." A victim of moderate hypothermia exhibiting the first seven signs listed below is still conscious but often confused. Severe hypothermia is determined by extreme skin coldness, loss of consciousness, faint pulse, and shallow, infrequent, or apparently absent respiration. Death is the ultimate result.

1. Severe shivering
2. Abnormal behavior
3. Slowing
4. Stumbling
5. Weakness
6. Inability to walk
7. Collapse
8. Stupor
9. Unconsciousness

Spill Containment

To prevent further spread of the anthrax spores:

- ④ Evacuate and secure the area.
- ④ Turn off the air handling systems, if possible.
- ④ Notify personnel to stay away from the site. Only those trained and informed about the release should access the impacted area.

Decontamination

Cleaning an area or item contaminated by anthrax involves numerous and variable issues that are specific to individual locations. No single technology, process, or strategy will be effective in every case. Responders must develop a [decontamination](#) plan that takes into account the:

- ④ Nature of the contamination.
- ④ Extent of the contamination.
- ④ Objectives of decontamination.



The extent of contamination and how the contamination was spread are critical considerations in isolating affected areas and selecting appropriate decontamination technologies.

Emergency Response

Notify the agencies below. Keep the contact numbers easily accessible.

- ④ Call **9-1-1** for local police assistance and the local Federal Bureau of Investigations (FBI).
- ④ Call the **National Response Center (NRC) at (800)424-8802**. The NRC will notify the appropriate agencies.



An emergency response protocol should be contained in the HASP for site personnel to follow while waiting and preparing for the notified agencies to arrive, initial actions should focus on the following:

- ④ Isolate contaminated areas,
- ④ Minimize exposure to others, and
- ④ Keep track of those who may have been exposed.

Standard Operating Procedures

Developing standard operating procedures (SOPs) provides a guideline for personnel to perform work activities as well as proper response as outlined in the HASP. Easily accessible SOPs can prevent mistakes and incorrect protocol, and therefore result in a safer workplace. Procedures directly related to an anthrax-related threat that may be included in the SOPs are:

- ④ [Notification procedures](#)
- ④ [Containment of the impacted area](#)
- ④ [Threat assessment](#)
- ④ [Evacuation procedures](#)

Additional information about evacuation procedures can be found in OSHA's [Evacuation Plans and Procedures eTool](#).

| [Home](#) | [What is Anthrax?](#) | [Who is at risk?](#) | [How do I prepare?](#) |
| [What if I have an anthrax threat?](#) | [How do I clean up my worksite?](#) |
| [Viewing/Printing Instructions](#) | [Credits](#) |

 [Back to Top](#)

www.osha.gov

www.dol.gov

[Contact Us](#) | [Freedom of Information Act](#) | [Information Quality](#) | [Customer Survey](#)
[Privacy and Security Statement](#) | [Disclaimers](#)



ANTHRAX e-Tool Protecting the Worksite against terrorism

- Home
- What is anthrax?
- Who is at risk?
- How do I prepare?
- What if I have a threat?
- How do I clean up my worksite?

What training requirements exist for workers involved in anthrax response and remediation?

A site-specific training program ensures that workers receive the hazard awareness training they need to work safely. The level of training required will depend on the types of activities that workers are performing. For anthrax response and remediation activities, training requirements are categorized as follows:



- ④ [Emergency Response](#)
- ④ [Cleanup Operations](#)
- ④ [Post-Emergency Cleanup](#)
- ④ [Appropriate Safety and Health Training](#)
- ④ [Anthrax-Specific Hazard Awareness Training](#)
- ④ [Additional Training Information](#)

Emergency Response

The five levels of training for employees who initially respond to an emergency are listed from the lowest to highest level of competency below:

- ④ **First Responder awareness level:** First responders at the awareness level are individuals who are likely to witness or discover a hazardous substance release and who have been trained to initiate an emergency response sequence by notifying the proper authorities of the release. They would take no further action beyond notifying the authorities of the release.
- ④ **First Responder operations level:** First responders at the operations level are individuals who respond to releases or potential releases of hazardous substances as part of the initial response to the site for the purpose of protecting nearby persons, property, or the environment from the effects of the release. They are trained to respond in a defensive fashion without actually trying to stop the release. Their function is to contain the release from a safe distance, keep it from spreading, and prevent exposures.



- ④ **Hazardous Materials Technician:** Hazardous

materials technicians are individuals who respond to releases or potential releases for the purpose of stopping the release. They assume a more aggressive role than a first responder at the operations level in that they will approach the point of release in order to plug, patch, or otherwise stop the release of a hazardous substance.



④ **Hazardous Materials Specialist:** Hazardous materials specialists are individuals who respond with and provide support to hazardous materials technicians. Their duties parallel those of hazardous materials technicians. However, those duties require a more directed or specific knowledge of the various substances they may be called upon to contain. The hazardous materials specialist would also act as the site liaison with federal, state, local, and other government authorities in regards to site activities.



④ **On-scene Incident Commander:** Incident commanders will assume control of the incident scene beyond the first responder awareness level.



Each level requires employers to have sufficient training or to have sufficient experience to objectively demonstrate competencies listed in [29 CFR 1910.120\(q\)\(6\)](#). Certification of training is required.

Cleanup Operations

At sites where OSHA's HAZWOPER standard applies, the safety and health training program should be based on the job hazard analysis in the [Health and Safety Plan](#) (HASP) and other relevant OSHA requirements. The training elements required by HAZWOPER include:

- ④ Initial anthrax hazard awareness training for site workers and supervisors,
- ④ Exceptions to initial training requirements,
- ④ Site-specific anthrax hazard awareness briefings for visitors and workers,
- ④ Refresher training,
- ④ Qualification of trainers,



- ④ Training certification, and
- ④ Emergency response training.

All employees who work on a HAZWOPER cleanup site (not limited to cleanup crew) where they are exposed to hazardous substances, health hazards, or safety hazards, must have training that meets the requirement of [29 CFR 1910.120\(e\)](#) or have equivalent experience and/or training. The four levels of training for employees who work on cleanup operations are listed below:

- ④ General site workers,
- ④ Workers on site only occasionally for a specific limited task (unlikely to be exposed over limits and not required to wear respirators),
- ④ Workers regularly on site in monitored and fully characterized task areas (unlikely to be exposed over limits and not required to wear respirators), and
- ④ Managers and supervisors.

Each level requires employees to have sufficient training or to have equivalent experience. Certification of training is required. The required elements of training are:

- ④ Names of personnel and alternates responsible for site safety and health,
- ④ Safety, health, and other hazards present on the site,
- ④ Use of personal protective equipment ([PPE](#)),
- ④ Work practices by which the employer can minimize risks from hazards,
- ④ Safe use of engineering controls and equipment on the site,
- ④ Medical surveillance requirements including recognition of the symptoms and signs that might indicate exposure to hazards,
- ④ Contents of the site safety and health plan including:
 - ④ Decontamination procedures in accordance with [29 CFR 1910.120\(k\)](#),
 - ④ An emergency response plan meeting the requirements of [29 CFR 1910.120\(l\)](#) for safe and effective responses to emergencies including the necessary PPE and other equipment,
 - ④ Confined space entry procedures, and
 - ④ A spill containment program meeting the requirements of [29 CFR 1910.120\(j\)](#).

Post-Emergency Cleanup

Where the cleanup is done on plant property using plant or workplace employees, these employees must have completed the training requirements of the following:

- ④ [Emergency action plan 29 CFR 1910.38\(a\)](#),
- ④ [Respiratory protection 29 CFR 1910.134](#),
- ④ [Hazard communication 29 CFR 1910.1200](#), and
- ④ Other appropriate safety and health training made necessary by the tasks that are expected to be performed (such as [PPE](#) and [decontamination](#) procedures).



Appropriate Safety and Health Training

A site-specific training program ensures that workers receive the training they need to work safely. Workers must receive all training required by applicable OSHA standards. This training may be included in the HAZWOPER curriculum. Examples of relevant training required by other standards include:

- ④ [Hazard communication](#)
- ④ [PPE](#)
- ④ [Respiratory protection](#)
- ④ [Fire extinguisher](#)
- ④ [Emergency action plan](#)
- ④ [Fire prevention plan](#)
- ④ [Emergency response](#)
- ④ [Lockout/tagout](#)
- ④ Observing working surfaces
- ④ [Noise](#)

Anthrax-Specific Hazard Awareness Training

Anthrax-specific hazard awareness training should help workers understand the health hazards of anthrax and how to protect themselves from exposure to spores. Specific topics might include:

- ④ How workers might be exposed to spores, the signs and [symptoms of infection](#), and medical conditions that could place them at increased risk (such as compromised immune systems),
- ④ Where contamination has been identified in the facility, and the status of [decontamination](#) of those areas, and
- ④ How to minimize the risk of disease through specific standard operating procedures and controls (such as engineering controls, work practices, housekeeping, or [PPE](#)), and whether specific measures are expected to be temporary or permanent.

There are additional training requirements for workers preparing contaminated materials or other hazardous materials for transportation to a treatment or disposal facility. These requirements can be found in the federal hazardous materials transportation regulations at [49 CFR Part 172, Subpart H](#).

Additional Training Information

- ④ [Code of Federal Regulations website](#)
- ④ Hazardous Waste Operations and Emergency Response (HAZWOPER), [29 CFR 1910.120 \[July 1, 2002\]](#) (the most current revision available at the time of this website's creation)

- Ⓒ Transportation Training, [49 CFR 172, Subpart H \[October 1, 2001\]](#) (the most current revision available at the time of this website's creation)
- Ⓒ Centers for Disease Control and Prevention (CDC), Morbidity and Mortality Weekly Report (MMWR), [Notice to Readers: Occupational Health Guidelines for Remediation Workers at Bacillus anthracis-Contaminated Sites \[September 6, 2002 / 51\(35\);786-789\]](#)

| [Home](#) | [What is Anthrax?](#) | [Who is at risk?](#) | [How do I prepare?](#) |
| [What if I have an anthrax threat?](#) | [How do I clean up my worksite?](#) |
| [Viewing/Printing Instructions](#) | [Credits](#) |

 [Back to Top](#)

www.osha.gov

www.dol.gov

[Contact Us](#) | [Freedom of Information Act](#) | [Information Quality](#) | [Customer Survey](#)
[Privacy and Security Statement](#) | [Disclaimers](#)

Occupational Safety & Health Administration
200 Constitution Avenue, NW
Washington, DC 20210



ANTHRAX e-TOOL Protecting the Worksite against terrorism

- Home
- What is anthrax?
- Who is at risk?
- How do I prepare?
- What if I have a threat?
- How do I clean up my worksite?

What types of Personal Protective Equipment (PPE) are necessary?

Personal protective equipment (PPE) shields or isolates workers from health and safety hazards in the workplace. In a site where anthrax spores may be present, PPE protects workers from exposure to respiratory and skin hazards and prevents the spread of contaminants to uncontaminated areas.



- ④ [Skin Protection](#)
- ④ [Respiratory Protection](#)
- ④ [PPE Training](#)

Skin Protection

Wearing protective clothing protects the skin and can prevent the transfer of contamination off-site. The appropriate level of skin protection, which should be outlined in the [HASP](#), is determined based on the site conditions. The appropriate level of skin protection used at an anthrax-impacted site should include, at a minimum, the following:

- ④ Tyvek or equivalent coveralls
- ④ Unpowdered disposable gloves made of lightweight nitrile or vinyl



Appropriate PPE selection will take into account the following:

- ④ Does it protect the skin from contact with anthrax spores?
- ④ Does it protect the worker from contact with chemicals used in the response?
- ④ Does it protect the worker from other site hazards identified in the HASP?
- ④ Can it be taken off safely and left at the site to prevent the transfer of contamination off-site.



The following links include more detailed information on Skin Protection:

- ④ [OSHA's PPE Technical Links page](#)

Respiratory Protection

Since airborne spores generally pose the greatest threat to personnel, respiratory protection is a necessary component of the PPE program. The OSHA respiratory protection standard ([29 CFR 1910.134](#)) requires you to establish and comply with an effective respiratory program. Program requirements include:



- ④ Program administration;
- ④ Worksite-specific procedures, respirator selection, employee training, fit testing;
- ④ Medical evaluation; and
- ④ Respirator use, cleaning, maintenance, and repair.

To date, experience has shown the following to provide adequate protection for most response, sampling, and decontamination activities involving spores:

- ④ Powered air-purifying respirators (PAPRs) with P100 filters, or
- ④ Full-face negative pressure air purifying respirators (APRs) with N95 filters.

Some emergency response operations may require that first responders wear the following with their normal Hazardous Material response ensembles:

- ④ Self-Contained Breathing Apparatus (SCBA).

The following links include more detailed information on Respiratory Protection:

- ④ [OSHA's Respiratory Protection Technical Links page](#)
- ④ [OSHA's PPE Technical Links page](#)
- ④ [OSHA Standard 29 CFR 1910.134, Respiratory Protection](#)

PPE Training

It is essential that personnel be trained on the proper use of PPE to ensure they receive greatest protection possible. Employers should certify in writing that the training has been provided and that employees understand what they need to know about PPE. The certification should show the name of each employee trained and the dates and types of training provided.

Workers should be trained to know:

- ④ When PPE is necessary
- ④ What type of PPE to use
- ④ Where the PPE is located
- ④ How the PPE should be worn



- ④ What the PPE limitations are
- ④ How long the PPE is likely to last
- ④ How to properly maintain and dispose of the PPE

In addition, appropriate personnel decontamination and contamination containment procedures are needed for workers using PPE to prevent exposure to anthrax. In general, these procedures are similar to those for asbestos abatement and include:

- ④ Isolating contaminated areas
- ④ Negative-pressure ventilation
- ④ A three- to five-stage decontamination line with a shower for equipment and personnel
- ④ Appropriate waste disposal

For additional detailed information about Respiratory Protection see the following:

- ④ [OSHA's Respiratory Protection Technical Links page](#)
- ④ [OSHA's PPE Technical Links page](#)
- ④ [Anthrax eTool Training Page](#)
- ④ EPA's publication "*Guidance for Controlling Asbestos-Containing Materials in Buildings*"; EPA-560/5-85-024 (Purple Book)
- ④ [OSHA's Hazardous Waste Operations and Emergency Response \(HAZWOPER\) Standard 29 CFR 1910.120](#)
- ④ [OSHA Standard 29 CFR 1910 Subpart I, Personal Protective Equipment](#)

[| Home |](#) [What is Anthrax? |](#) [Who is at risk? |](#) [How do I prepare? |](#)
[What if I have an anthrax threat? |](#) [How do I clean up my worksite? |](#)
[| Viewing/Printing Instructions |](#) [Credits |](#)

 [Back to Top](#)

www.osha.gov

www.dol.gov

[Contact Us |](#) [Freedom of Information Act |](#) [Information Quality |](#) [Customer Survey](#)
[Privacy and Security Statement |](#) [Disclaimers](#)

Occupational Safety & Health Administration
200 Constitution Avenue, NW
Washington, DC 20210



ANTHRAX e-TOOL Protecting the Worksite against terrorism

- Home
- What is anthrax?
- Who is at risk?
- How do I prepare?
- What if I have a threat?
- How do I clean up my worksite?

What are the employee medical program requirements?

Because of the potential for exposure to anthrax, medical measures to prevent anthrax must be taken. The medical measures differ depending on the type of worker and duration of exposure. There are three types or categories of workers or others who may be exposed to anthrax:



- Ⓞ [Short-Term Response Workers](#)
- Ⓞ [Long-Term Response Workers](#)
- Ⓞ [Occupants, Workers, or Visitors at a Site Contaminated with Anthrax](#)

In addition to the medical measures to prevent exposure to anthrax, and the medical surveillance programs required in the HAZWOPER ([29 CFR 1910.120](#)) and respiratory protection regulations, the occupational health and safety plan ([HASP](#)) should also include the following two major components to protect workers:

- Ⓞ [Medical Screening and Follow-Up Care for Anthrax and Medical Complications Related to Preventive Measures](#)
- Ⓞ [Knowledge and Information that Workers Need to Prevent Anthrax and Medical Complications Related to Preventive Measures](#)

Additional information regarding medical surveillance can be found at the following site:

- Ⓞ [Centers for Disease Control and Prevention \(CDC\), Morbidity and Mortality Weekly Report \(MMWR\), Notice to Readers: Occupational Health Guidelines for Remediation Workers at Bacillus anthracis-Contaminated Sites \[September 6, 2002 / 51\(35\);786-789\]](#)

Short-Term Response Workers

Exposures are limited to a single episode or a few episodes within a brief period (less than 30 days). Local emergency medical personnel, police, and firefighters who are not expected to re-enter contaminated areas for longer periods of time fall into this category.



Short-term response workers should be:

- Ⓞ Adequately immunized with anthrax vaccine, or
- Ⓞ Placed on appropriate prophylactic antibiotics at the time of their first exposure and continue for 60 days after their last exposure. Because experimental data indicate that viable spores may persist in the lungs for 100 days after exposure, an option is the use of antibiotics for 100 days after exposure.

Long-Term Response Workers

These individuals have repeated exposures over longer periods of time (30 days or more). Environmental response team members and decontamination workers fall into this category. They may work at multiple sites (such as industrial hygienists conducting environmental sampling) or at a single site (such as contractors performing decontamination work).



Long-term response workers should be:

- ④ Adequately immunized with anthrax vaccine before exposure if they are at high risk of repeated exposures.
- ④ Placed on appropriate prophylactic antibiotics, if necessary, for considerably longer than the 60 days recommended for short-term workers. Because experimental data indicate that viable spores may persist in the lungs for 100 days after exposure, an option is the use of antibiotics for 100 days after exposure.
- ④ If the vaccine series is started after exposure has occurred, antibiotic prophylaxis should be continued during the first three doses to provide protection until an adequate immune response has developed.

Occupants, Workers, or Visitors at a Site Contaminated with Anthrax

Postal or office workers or maintenance and housekeeping personnel would fall into this category. The medical program for this group should have two phases: the immediate post-exposure period and the period after a previously contaminated site has been cleared for unrestricted entry and occupancy. Initial medical screening should be done to identify exposed persons who should avoid taking antibiotics.



Medical measures during the immediate post-exposure period:

- ④ Individuals should be placed on appropriate prophylactic antibiotics at the time of their first exposure and continue for 60 days after their last exposure. Because experimental data indicate that viable spores may persist in the lungs for 100 days after exposure, an option is the use of antibiotics for 100 days after exposure.

Once a previously contaminated site has been cleared for re-occupancy, personal protection and medical measures to prevent anthrax are presumably no longer necessary. However, a precautionary program of medical monitoring may be prudent to assure that anthrax is no longer a threat. The program should be designed and administered under the supervision of a licensed physician.

Medical measures after a previously contaminated site has been cleared for unrestricted entry and occupancy should include:

- ④ Initial medical history to screen for high-risk conditions (such as compromised immunity, skin conditions)
- ④ Counseling of high-risk persons
- ④ Around-the-clock access to medical coverage for anthrax-like symptoms
- ④ Confidentiality of medical information

Medical screening and follow-up care for anthrax and medical complications related to preventive measures

The purpose of occupational health surveillance in the workplace is to improve the effectiveness of the occupational health and safety program by systematically collecting and analyzing information that pertains to at-risk workers.

The program for monitoring exposed persons includes:

④ **Medical screening program**

The medical screening program is the use of examinations or tests to detect adverse effects on a worker's health at an early stage when prevention is possible or treatment is most effective. It should include:

- ④ Baseline medical screening to identify pre-existing conditions that may affect an individual worker's fitness for duty, and who should avoid antibiotics or vaccines.
- ④ Periodic evaluations to reassess fitness for duty and to detect symptoms of the development of anthrax or adverse effects related to preventive measures (such as antibiotics).
- ④ Final evaluation when it is no longer necessary for a worker to re-enter a contaminated site, to identify changes from the baseline and any new risk factors.



④ **Medical monitoring program for adverse work-related health effects**

The medical monitoring program for adverse effects related to antibiotic use should include the following so that an informed decision appropriate for the affected individual can be made:

- ④ Plans to inform affected workers about available options for preventing anthrax
- ④ Risks and benefits of each option

④ **Exposure Monitoring (such as incidents of breaches in PPE)**

There are no validated methods for monitoring a person's exposure to *Bacillus anthracis*. There are useful epidemiologic tools that can be used, but these should not be used to assess a person's exposure or to make decisions about the use of antibiotics or vaccine. These tools include:

- ④ Nasal swabs
- ④ Serology

Inhalation exposure to a high concentration of *Bacillus anthracis* spores may rapidly result in death. Therefore, exposure to aerosolized powder known or highly suspected to be contaminated with *Bacillus anthracis* spores should be treated as a medical emergency.

**Knowledge and Information that Workers Need to Prevent
Anthrax and Medical Complications Related to Preventive Measures**

In addition to HAZWOPER training ([29 CFR 1910.120](#)), response workers will also need additional knowledge and information about anthrax and the medical measures that can protect them. It should be stressed that for preventive measures to be effective, individual workers must fully understand them and use them correctly. All response workers should receive training to:

- ④ Recognize and report early symptoms and signs of anthrax,
- ④ Understand the importance of immediate medical attention,
- ④ Know how to access emergency medical care,
- ④ Know about potential adverse effects and interactions with food and drugs if taking antibiotics, and

- ④ Understand the potential adverse effects of vaccine and the amount of time necessary to develop an immune response if using the vaccine as a preventive measure.

The validity and reliability of symptom reports are only as good as the exposed person's knowledge and understanding of the characteristics of anthrax and risks for developing the disease. Successful treatment of anthrax will depend on the individual's understanding of the need for immediate medical attention should symptoms occur and knowledge of how to access emergency medical care. Therefore, hazard awareness training is an important component of the medical program.

The following links include more detailed information on Medical Program Requirements:

- ④ [OSHA's Hazardous Waste Operations and Emergency Response \(HAZWOPER\) Standard 29 CFR 1910.120](#)
- ④ [Anthrax eTool Disease Recognition page](#)

| [Home](#) | [What is Anthrax?](#) | [Who is at risk?](#) | [How do I prepare?](#) |
| [What if I have an anthrax threat?](#) | [How do I clean up my worksite?](#) |
| [Viewing/Printing Instructions](#) | [Credits](#) |

 [Back to Top](#)

www.osha.gov

www.dol.gov

[Contact Us](#) | [Freedom of Information Act](#) | [Information Quality](#) | [Customer Survey](#)
[Privacy and Security Statement](#) | [Disclaimers](#)

Occupational Safety & Health Administration
200 Constitution Avenue, NW
Washington, DC 20210



ANTHRAX e-Tool Protecting the Worksite against terrorism

- Home
- What is anthrax?
- Who is at risk?
- How do I prepare?
- What if I have a threat?
- How do I clean up my worksite?

How do I sample and analyze for anthrax?

Environmental sampling is an important tool for determining the presence of *Bacillus anthracis* spores in indoor environments. Sampling can help assess the extent and degree of contamination and the risk of exposure to building occupants and responders. Sample results also contribute to informed decisions on medical treatment and decontamination options and are ultimately used to determine the effectiveness of decontamination.



- ④ [General Sampling Plan](#)
- ④ [Specific Sampling Objectives](#)
- ④ [Sampling Approach](#)
- ④ [Sampling Techniques](#)
- ④ [Analytical Methods](#)
- ④ [Packaging and Transportation of Samples](#)
- ④ [Interpretation of Results](#)
- ④ [Coordination with Affected Parties](#)

General Sampling Plan


The purpose of a sampling plan is to create a road map to achieving defined sampling objectives. An effective sampling plan will provide confidence that the results obtained are valid and indicative of the contamination present. The sampling plan should address each of the following questions:

- ④ What are the [specific sampling objectives](#)?
- ④ What [sampling approach](#) will be used to meet the objectives?
- ④ What [analytical methods](#) will be used to test for contamination?
- ④ How will samples be [packed, handled, and transported](#)?

In evaluating methods that will achieve the sampling objectives, several other factors should be considered:

- ④ Laboratory capability (such as the laboratory's ability to handle the proposed sampling media),
- ④ Collection efficiency of the method, number of samples, and need for quantification,
- ④ Suitability of the sample-collection and analytical methods,
- ④ Cost effectiveness and efficiency of the sampling plan in meeting stated objectives, and
- ④ Utility of the sampling method to the owner and/or federal

agency having jurisdiction over the project.



Specific Sampling Objectives

There are a variety of potential sampling objectives. The specific sampling objectives that may be applicable for an anthrax response may include:

- ④ **Real-time Monitoring:** Determine, in real-time, whether a release of spores is occurring or has occurred in a facility. Real-time instruments may be used to detect biological agents as they are released.
- ④ **Preliminary Assessment of a Facility (Screening):** Determine qualitatively whether any spores are present. Typically, composite samples of large areas and air volumes are obtained to maximize the likelihood of finding contamination.
- ④ **Identification of Spores in a Bulk Material:** Determine qualitatively if a bulk material, such as a powder in an envelope, is contaminated with anthrax. On-site analysis may be used for preliminary assessment, but laboratory analysis provides confirmation.
- ④ **Determination of Contamination of an Article:** Determine whether the surface of a small article is contaminated. Typically, composite surface samples of large articles and/or individual samples of small articles are collected.
- ④ **Extent and Location of Contamination (Site Characterization):** After anthrax is positively identified, further sampling is necessary to determine how far the contamination has spread. Sampling is performed to determine qualitatively, and if possible, semi-quantitatively, the extent and magnitude of contamination.
- ④ **Effectiveness of Decontamination (Process Verification Sampling):** Determine whether decontamination has reduced spores to a safe level.
- ④ **Post-Decontamination Sampling (Re-occupancy Verification Sampling):** Final post-decontamination sampling is conducted inside and outside of the exclusion zone to verify that the originally contaminated environment has been sufficiently decontaminated to allow re-occupancy of the area without the use of [personal protective equipment](#) (PPE).
- ④ **Special Re-occupancy Considerations:** There may be site-specific circumstances where additional sampling should be considered.

Sampling Approach

After the sampling objectives have been identified, a logical approach and schedule must be developed to carry out the sampling tasks. The approach and schedule will depend on the objectives.

For Assessment/Characterization:

- ④ **Targeted Sampling Strategy:** If a building or area becomes contaminated from a known source and the source is quickly isolated, the sampling approach will center on the source area and move outward away from the source to define the extent of contamination.
- ④ **Statistically-Based Strategy:** If contamination is likely to be present in a building or area and the source has not been identified, sampling will focus on areas that are statistically the most likely to be contaminated. The objective of statistical sampling is to maximize the probability of detecting contamination.
- ④ **Air Movement and HVAC Considerations:** Spore-bearing particles less than 10 microns in size, or spores themselves, may remain suspended in the air for long periods of time (such as hours or days). In such cases, spores may spread throughout an air space and into adjacent areas on localized air currents, such as those created by people walking by, and also through generalized airflow created by HVAC systems. To fully assess the extent of contamination, the investigator should extend beyond the targeted approach to sample areas on projected contaminant pathways, such as those associated with air movement, dust collection, or work process flow.

For Verification Sampling Prior to Occupancy:

- ④ **Post-Decontamination Surface Sampling:** The effectiveness of decontamination should be confirmed by post-decontamination environmental sampling in ambient air; and in areas and on surfaces that were previously contaminated. Verification sampling prior to occupancy should include surfaces and air in areas outside of the exclusion zone (for example, the area where decontamination activities were conducted) to ensure that the outside areas continue to remain free of spores.
- ④ **Air Sampling:** Aggressive air sampling techniques have been developed for recent anthrax responses that model EPA guidance for clearing facilities for re-occupancy after asbestos decontamination. While the area is under negative pressure, all surfaces are aggressively agitated and air is continuously disturbed while samples are collected. An air sampling method that maximizes the likelihood of detecting contamination should be used.

Sampling Quality Assurance and Quality Control (QA/QC) Considerations:

QA/QC measures must be incorporated into the sampling approach to insure the legitimacy of sampling results. The sampling objectives will dictate the rigor of the QA/QC program for a given site or task.

QA/QC includes four key elements:

- ④ QC of field activities
- ④ Sample documentation and management
- ④ Sample handling and shipment
- ④ Data validation and management

Sampling Techniques

There are various sampling techniques available for determining the presence of anthrax in air and surface samples.

The following techniques are available and have proven useful for **surface and bulk sampling**:

- ④ Wet wipe,
- ④ Wet swab,

- ④ High-volume vacuum with "HEPA" sock (Alsock), and
- ④ Bulk sampling.



The following techniques are available and have proven useful for **air sampling**:

- ④ Gelatin filter (low volume),
- ④ Mixed cellulose ester (MCE) filter,
- ④ Anderson air sampler and single-stage impactors with settle plates,
- ④ Open agar plate,
- ④ Dry filter unit (high-volume air sampler with polyester 1-micron filter), and
- ④ Liquid impingers.



The following links provide additional information regarding sampling techniques:

- ④ [Bacillus Anthracis, Environmental Sampling on Surfaces](#). OSHA (2003, March), 13 pages. This document describes sampling and analysis methods for anthrax surface samples.
- ④ [Bacillus Anthracis Spores \(Etiologic Agent of Anthrax\) in Air](#). OSHA (2003, March), 22 pages. This document describes sampling and analysis methods for anthrax air samples.

Analytical Methods

The following analytical methods are available for the detection of *Bacillus anthracis*:

- ④ **Immunoassay Tests:** Hand-held assays, sometimes referred to as "smart tickets," are sold commercially for the rapid detection of *Bacillus anthracis* and other biological agents. These assays are intended only for screening environmental samples. Due to their low and questionable sensitivity, they cannot be relied on for a determination that anthrax is or is not present.
- ④ **Polymerase Chain Reaction (PCR):** PCR is a technique that amplifies DNA and compares sequences to known test probe standards for *Bacillus anthracis*. PCR can be used in the field or the laboratory; in either case, the samples must be cultured to confirm that the bacteria is *Bacillus anthracis* and that it is viable. Field PCR systems are very selective, but do not work well with heterogeneous environmental samples, and the probes are very expensive. PCR has been shown to work best as a final confirmation of positive samples taken from plated colonies.
- ④ **Culturing:** Samples may be analyzed for *Bacillus anthracis* using a traditional lab culturing technique. The sample is appropriately prepared for elution and plating, after which it is inoculated onto plates containing sheep blood agar. The plates are allowed to incubate for several hours and are then examined for growth of suspicious colonies.



The laboratory selected for the sample analyses should be consulted by the team developing the sampling plan. The lab should be authorized for work with *Bacillus anthracis* and laboratory procedures must conform to guidance provided by CDC or the Association of Public Health Laboratories (APHL). The lab should also be a part of the Laboratory Response Network (LRN), which was developed by CDC, APHL, and the FBI to help public health labs prepare for and respond to acts of terrorism, including bioterrorism.

The following links provide additional information regarding analytical methods:

- ④ [Bacillus Anthracis, Environmental Sampling on Surfaces](#). OSHA (2003, March), 13 pages. This document describes sampling and analysis methods for anthrax surface samples.
- ④ [Bacillus Anthracis Spores \(Etiologic Agent of Anthrax\) in Air](#). OSHA (2003, March), 22 pages. This document describes sampling and analysis methods for anthrax air samples.
- ④ Centers for Disease Control and Prevention, Public Health Emergency Preparedness and Response, [Approved Tests for the Detection of *Bacillus anthracis* in the Laboratory Response Network \(LRN\)](#)
- ④ Centers for Disease Control and Prevention, Public Health Emergency Preparedness and Response, [The Laboratory Response Network for Bioterrorism \(LRN\)](#) [a slideshow that provides a general overview of the LRN]

Packaging and Transportation of Samples

There are strict requirements for packaging and transporting anthrax samples to ensure that the general public and workers transporting the samples are protected from exposure. These requirements include:

- ④ Rigorous packaging designed to withstand rough handling and prevent leakage,
- ④ Appropriate marking and labeling that identifies the contents of the package,
- ④ Documentation of the hazardous contents of the package and emergency point-of-contact, and
- ④ Training of transportation workers on how to handle the contents in the event of an emergency.



Packaging and transporting anthrax samples are subject to various regulations established by the DOT, CDC, USPS, OSHA, and the International Air Transport Association (IATA). It is also important to consult with the analytical laboratory receiving the samples to determine whether they have additional packaging or shipping requirements.

Additional details on packaging and shipping procedures for anthrax samples, according to CDC guidelines, can be found at:

- ④ Centers for Disease Control and Prevention, Public Health Emergency Preparedness and Response, [Packing Critical Biological Agents](#) (protocols for packing, shipping, and transport of biological agents/diseases)
- ④ Centers for Disease Control and Prevention, Office of Health and Safety, Biosafety Branch, [Interstate Shipment of Etiologic Agents \(42 CFR Part 72\)](#)

Interpretation of Results

Since analytical methods are not fully validated for *Bacillus anthracis*, the following considerations are

recommended for interpretation of analytical results:

- ④ Use a multi-disciplinary team of technical experts to interpret the analytical results.
- ④ Consult with field investigators and laboratory personnel during the interpretation process. This will provide the best insight into sample collection and recovery.
- ④ Consider the analytical method limitations and use professional judgment in interpreting any positive or negative findings as well as quantitative or semi-quantitative results.
- ④ In the case of preliminary assessment sampling, positive findings will usually require more extensive sampling in that particular area.
- ④ Laboratories will sometimes report "colony counts" with positive results. In some cases, colony counts can help the data interpretation teams decide how much more extensively to sample in that area, prior to establishing additional isolation and starting decontamination. However, quantitative methods have not been validated for a variety of sample collection methods and the use of "colony count" data is not yet accepted as a reliable indicator of large populations of colony forming units.
- ④ The multi-disciplinary team described above should establish a decision tree for evaluating the adequacy of the sampling and decontamination to coordinate its efforts in reaching a unified conclusion.



Coordination with Affected Parties

Before, during, and after a response, the sampling team should coordinate with the following affected parties:

- ④ Facility manager,
- ④ Union officials, and
- ④ Other individuals who might have site-specific knowledge of the contamination.

The sampling plan should indicate the following:

- ④ Employees working within the affected sampling area need to be informed of the sampling objectives and methods.
- ④ The method of providing results to employees working within the affected sampling area.
- ④ Before sampling begins, employees should be told when, where, and why sampling may occur and they should also be advised when plans are changed.

[| Home](#) | [What is Anthrax?](#) | [Who is at risk?](#) | [How do I prepare?](#) |
[What if I have an anthrax threat?](#) | [How do I clean up my worksite?](#) |
[Viewing/Printing Instructions](#) | [Credits](#) |



ANTHRAX eTool Protecting the Worksite against terrorism

- [Home](#)
- [What is anthrax?](#)
- [Who is at risk?](#)
- [How do I prepare?](#)
- [What if I have a threat?](#)
- [How do I clean up my worksite?](#)

How should I decontaminate during response actions?

This page provides information on decontaminating buildings or specific areas, systems, or items within buildings after an actual release of anthrax. Anthrax decontamination is a repetitive process that may involve the use of multiple decontamination processes and technologies. Selection of appropriate technologies varies depending on factors specific to the release and to the technologies themselves, but the primary considerations are always the effectiveness and safety of the products and processes.



- ④ [Planning for Decontamination](#)
- ④ [Preparing for Decontamination](#)
- ④ [Decontamination Technologies](#)
- ④ [Judging the Effectiveness of Decontamination](#)

Planning for Decontamination

Cleaning an area or item contaminated by anthrax involves numerous and variable issues that are specific to individual locations. No single technology, process, or strategy will be effective in every case. Responders must develop a decontamination plan that takes into account the following:

- ④ **The nature of the contamination:** including the type of anthrax involved, how it entered the facility, and the physical characteristics that affect the spread of contamination.
- ④ **The extent of contamination:** including the amount of contamination and possible pathways by which it could have or will spread.
- ④ **The objectives of decontamination:** including the intended re-use of the facility and building systems and whether items will be decontaminated for re-use or treated for disposal.

The extent of contamination and how the contamination spread are critical considerations in isolating affected areas and selecting appropriate decontamination technologies. For example, if spores are widely dispersed and have traveled through the air, decontamination may involve extensive isolation and fumigation. In contrast, if the contamination is limited to a small area and spores are not likely to become airborne, then minimal isolation and surface decontamination methods alone may suffice.

All stakeholders--local authorities, building owners and residents, federal, state, and local environmental and health agencies, the affected public, and others--should be consulted before decontamination begins. A site health and safety plan (HASP) is needed to protect workers inside and outside the contaminated area, as well as the surrounding population. The facility manager should notify employees and others (such as union representatives) of the nature and scope of the work and its expected duration.

More detailed information about HASPs can be found on the [HASP](#) page of this eTool.

Preparing for Decontamination

The results of the sampling for extent of contamination should make it possible to distinguish between contaminated and uncontaminated areas and to determine the types of surfaces involved. To prevent the spread of contamination by movement of workers or equipment, it may be advisable to isolate the contaminated area, depending on the impacted area and the extent of contamination.

Decontamination should address the following:

- ④ **Hidden sources of contamination:** Desktop computers and other objects with internal fans that draw air into the case may have filters or electrostatic devices to control dust intake. These filters or the equipment chassis may be a reservoir of contamination. If selected technologies may damage the item or may not penetrate to hidden locations, then these items may be dealt with in an alternative manner. The manufacturer of the device should be consulted if it is to be saved for re-use.
- ④ **Pre-cleaning:** Excessive amounts of dirt or other organic material on the surface to be cleaned may decrease the effectiveness of the selected decontamination method. Using certain techniques, such as [HEPA vacuuming](#), to remove some of the dirt and debris could reduce the need to perform more aggressive chemical decontamination.
- ④ **Removal of items:** To reduce potential spread of contamination, items should be decontaminated in place. If the selected technology will destroy an item that must be salvaged, then the item may be removed and decontaminated elsewhere with an alternative technology. This requires a means of safe transport and a separate isolation chamber, which adds complexity to the decontamination process.



Additional, more detailed information about sampling for *Bacillus anthracis*, can be found on the [Sampling](#) page of this eTool.

Decontamination Technologies

Decontamination technologies can be divided into three categories:

1. **Surface decontamination products**, which are used to treat spores on hard, non-porous surfaces such as desks, walls, and hard flooring. There are two methods to treat surfaces:

- ④ **High Efficiency Particulate Air (HEPA) Vacuuming:** accomplishes two purposes: (1) it helps remove dirt and other debris that may reduce the effectiveness of subsequent decontamination, and (2) it also removes some of the spores, reducing the number that must be killed by subsequent decontamination. An advantage of this technology is that there is little potential for damage to furnishings. A limitation of this technology is that it can only remove surface contamination. The operator must also avoid allowing the exhaust to stir the air in the affected room and must safely dispose of contaminated filters.
- ④ **Liquid Antimicrobial Products - Not for Porous Surfaces:** may be used to inactivate spores on hard surfaces only. These products--which can be applied by pouring, mopping, or spraying--include oxidizing, bleaching, or other agents such



HEPA filters can be attached to high-volume vacuums such as this.

as aqueous chlorine dioxide, sodium hypochlorite, hydrogen peroxide, and peroxyacetic acid combined. Several factors should be considered when deciding which liquid antimicrobial products to use and how to apply them. Each product affects surfaces differently in terms of corrosivity, staining, and residue. These products will be effective only if the directions for use of the product are followed precisely (such as mixing directions, application method and dosage rate, pre-cleaning of surfaces, and contact time).



2. **Fumigation**, which involves use of an antimicrobial gas or vapor to destroy aerosolized spores and spores adhered to non-porous and porous surfaces. In addition to decontaminating a variety of surfaces, fumigants are able to decontaminate airborne spores that a surface cleaner would miss.
3. **Other decontamination products**, which are primarily used in chambers or other specialized equipment. Technologies that can be used to decontaminate specific items outside the affected area or environment include the following:
 - ④ **Chemical Sterilization:** chemicals such as ethylene oxide, chlorine dioxide, or paraformaldehyde are used to kill spores on discrete items placed in a sterilization chamber. Sufficient aeration of the items following treatment is necessary to remove residual amounts of the sterilant and any toxic by-products that may have formed. For effective decontamination, each chemical sterilant has specified ranges of temperature, relative humidity, concentration, and duration of application.
 - ④ **Irradiation:** irradiation techniques, including cobalt-60 and electron beam technologies, can effectively destroy anthrax. These techniques are generally available only for off-site decontamination. They may destroy magnetic media such as film or videotape, and they tend to be expensive.

Selection of the appropriate technology will require an evaluation of the specific site conditions and nature of contamination. Other considerations include the conditions required for effective application (for example, humidity for fumigations or pH for certain surface treatments), how the technology will affect the area or item being treated, and the risks associated with use (such as physical, chemical, and toxicologic parameters of the product).

The Environmental Protection Agency's (EPA) Technology Innovation Office (TIO) has developed an information clearinghouse (www.EPATechBIT.org) as a centralized location to collect and disseminate information about decontamination technologies and also for technology vendors to provide information.

Judging the Effectiveness of Decontamination

There are separate criteria, described below, for judging the effectiveness of decontamination of objects in an off-site sterilization chamber, and for decontamination of sites such as offices or buildings.

- ④ For objects decontaminated in an off-site sterilization chamber, biological indicators such as surrogate spore test strips, may be placed in the chamber along with the objects. Although the optimal results of decontamination would be if the biological indicators showed no evidence of bacterial growth, OSHA believes that there may be safe alternatives to the "no growth" decontamination.
- ④ To determine whether decontamination of a site has been effective, a rigorous round of environmental sampling should be performed following the decontamination process, and the samples should be cultured for *Bacillus anthracis* in a nationally accredited lab. Rigorous environmental sampling should be done in all decontaminated sites, regardless of type of technology used or the extent of the decontamination. In areas that have been fumigated,

biological indicators (such as surrogate spore strips) may be used to determine whether the fumigant has effectively permeated the area under specific conditions (such as concentration, time, temperature, and relative humidity) sufficient to kill *Bacillus anthracis* spores. The results of the culture of both the environmental samples and the biological indicators should be evaluated to determine the effectiveness of the fumigation. Again, the optimal results of decontamination would be if the biological indicators showed no evidence of bacterial growth. However, OSHA believes that there may be safe alternatives to the "no growth" decontamination.

A different technology may be used for further decontaminating an area in which sampling showed the presence of viable anthrax spores. For example, if evidence of bacterial growth is found only on a desk in an office that was fumigated, an approved liquid sporicidal product may be applied to complete decontamination of the desk.

Once an area has been determined to be effectively decontaminated, there is no guarantee that all viable spores have been eliminated, even when post-decontamination samples show no growth. Moreover, statistical calculations of the effectiveness of sampling and analytical methods indicate that some spores may remain in a decontaminated area even though environmental sampling shows "no growth." Nevertheless, the potential risk of a person contracting the disease in such an area is considered to be extremely low.

Although the "no growth" decontamination goal may be the best possible way to ensure safety in re-occupying a decontaminated area, OSHA believes that there may be safe alternatives, especially in workplace situations where the use of PPE, special work practices, or other engineering controls might also minimize the risk of disease.

This approach is consistent with the NCP, which allows case-by-case decisions based on the future use of the site and other controls that might be used to ensure safety. Decisions on alternative options such as these would have to be site-specific, and supported by experts in epidemiology, public health, industrial hygiene, and environmental protection. The basis for these types of decisions should also be thoroughly documented.

Additional information about decontamination techniques and process, and post-decontamination sampling strategies can be found on the [Sampling](#) page of this eTool.

| [Home](#) | [What is Anthrax?](#) | [Who is at risk?](#) | [How do I prepare?](#) |
| [What if I have an anthrax threat?](#) | [How do I clean up my worksite?](#) |
| [Viewing/Printing Instructions](#) | [Credits](#) |

 [Back to Top](#)

www.osha.gov

www.dol.gov

[Contact Us](#) | [Freedom of Information Act](#) | [Information Quality](#) | [Customer Survey](#)
[Privacy and Security Statement](#) | [Disclaimers](#)

Occupational Safety & Health Administration
200 Constitution Avenue, NW
Washington, DC 20210



ANTHRAX e-TOOL Protecting the Worksite against terrorism

- Home
- What is anthrax?
- Who is at risk?
- How do I prepare?
- What if I have a threat?
- How do I clean up my worksite?

What is a transition program and what elements should I include in my transition program?

A transition program is required for facilities in which anthrax contamination has been identified and normal work operations are conducted. Three examples of this situation include the following:



1. Normal work activities were resumed in a contaminated facility following decontamination and clearance sampling efforts.
2. Original sampling identified low levels of surface and air contamination and therefore occupancy is permitted.
3. Occupancy is permitted outside an established exclusion zone in which decontamination is being performed.

The duration of the program would generally be dictated by sampling results obtained during the transitional period. Site-specific transitional elements should be developed for each facility and incorporated as part of the regular site health and safety plan (HASP). It is recommended that an industrial hygienist develop these elements or at a minimum provide input during the development phase. Transitional Program elements to be included in the HASP include some or all of the following elements, as applicable:

- ④ [Hazard Awareness Training](#)
- ④ [Medical Surveillance](#)
- ④ [Transitional Sampling](#)
- ④ [Personal Protective Equipment](#)
- ④ [Personal Hygiene](#)
- ④ [Interim Standard Operating Procedures](#)

Hazard Awareness Training

Hazard awareness training is intended to communicate information concerning hazards of anthrax and appropriate protective measures to employees. This program may include, but is not limited to:

- ④ Elements of the transition program.
- ④ The health hazards of anthrax, including routes of entry, signs and [symptoms of exposure](#), synergistic effects, and any medical conditions which would place employees at increased risk (i.e., immunocompromised individuals).
- ④ Operations in the work area where anthrax has been identified.
- ④ Dissemination of sampling results to employees, including information on how to gain access to such results.
- ④ Any applicable control measures, such as appropriate engineering controls, work practices,



housekeeping, or [personal protective equipment](#) (PPE).

- ④ Implementation of interim standard operating procedures to control anthrax exposure during operations, maintenance, cleaning, etc.
- ④ Frequent updates regarding any on-going anthrax sampling, decontamination, control, medical surveillance, and related activities being performed at the facility, as applicable.

Additional detailed information may be found at the following site:

- ④ [Anthrax eTool Training Page](#)

Medical Surveillance

A medical surveillance program is necessary to ensure that employees receive appropriate preventive care. Medical surveillance includes, but is not limited to:

- ④ Identification of employee population at risk and establishment of controls for such employees (such as work reassignment, [PPE](#), prophylactic medication, etc.).
- ④ Administrative follow-up on absentees (such as sick leave, etc.).
- ④ Selection of prophylactic medication, as appropriate.
- ④ Response to symptoms reported by employees.



Additional detailed information may be found at the following sites:

- ④ [OSHA's Hazardous Waste Operations and Emergency Response \(HAZWOPER\) Standard 29 CFR 1910.120](#)
- ④ [Anthrax eTool Disease Recognition page](#)
- ④ [Anthrax eTool Medical Program page](#)

Transitional Sampling

Transitional sampling is conducted to confirm that the occupied areas remain safe for occupancy. Sampling during this period is continued until repeatable results demonstrate that contamination remains below an established target level. Additional information can also be found in the chapter on sampling. Elements of transitional sampling include, but are not limited to, the following:

- ④ Determination of appropriate sampling techniques. Recommended techniques include non-aggressive high-volume air sampling, aggressive HEPA vacuum surface sampling, and if appropriate bulk sampling (such as bulk samples from HEPA vacuum bags used to clean surfaces).
- ④ Use of high-volume air sampling as an essential tool in order to characterize levels of anthrax in the air and provide inhalational exposure information to employees.
- ④ Identification of specific locations and frequency of

sampling.



Additional detailed information may be found at the following site:

- ④ [Anthrax eTool Sampling page](#)

Personal Protective Equipment

The workplace must be reassessed in order to select and use appropriate [personal protective equipment](#) (PPE) to protect employees from the anthrax hazards present. The specific types of PPE utilized would be dependant on the actual operation in question and results from the reassessment. Examples of work operations where modifications to PPE may be necessary are as follows:



- ④ Operating equipment or working on surfaces where anthrax was previously identified.
- ④ Performing maintenance tasks, such as cleaning equipment or changing HEPA vacuum filters.

Additional detailed information may be found at the following site:

- ④ [OSHA's PPE Technical Links page](#)

Personal Hygiene

A personal hygiene program should be implemented to limit exposures and control the spread of anthrax contamination. Procedures that may be required include the following:

- ④ Assuring that food or beverage is not present or consumed, tobacco products are not present or used, and cosmetics are not applied.
- ④ Regular washing of the hands and/or face, and before eating, drinking, using tobacco, or applying cosmetics.
- ④ Showering as necessary.



Interim Standard Operating Procedures

Interim standard operating procedures (SOPs) must be developed to address special work activities necessary under the transitional program. Affected employees should receive training on the interim SOPs. These SOPs may include, but are not limited to, the following topics:

- ④ Maintenance and housekeeping procedures developed or modified to control the spread of contamination and protect employees. Examples of procedures include:
 - ④ Use of HEPA vacuums for cleaning surfaces instead of sweeping or other methods,
 - ④ Cleaning, maintenance, and filter and bag removal for HEPA vacuums,
 - ④ Maintenance and cleaning of facility equipment,
 - ④ Cleaning of floors and other surfaces, and
 - ④ Handling and disposal of wastes.

- ④ Changes to regular work operations and equipment, as applicable.

- ④ Modifications to facility-wide mechanical systems, particularly heating, ventilation, and air-conditioning (HVAC) systems. Examples of HVAC modifications include:
 - ④ Increase in ventilation rates (air changes per hour)
 - ④ Increase in percentage of outside air
 - ④ Use of HEPA filters to collect dust in circulated air

- ④ Other applicable major elements being implemented as part of the transitional program, as described previously (such as training, medical surveillance, sampling, PPE, and hygiene).



Additional detailed information may be found at the following site:

- ④ [Anthrax eTool HASP page](#)

| [Home](#) | [What is Anthrax?](#) | [Who is at risk?](#) | [How do I prepare?](#) |
| [What if I have an anthrax threat?](#) | [How do I clean up my worksite?](#) |
| [Viewing/Printing Instructions](#) | [Credits](#) |

 [Back to Top](#)

www.osha.gov

www.dol.gov

[Contact Us](#) | [Freedom of Information Act](#) | [Information Quality](#) | [Customer Survey](#)
[Privacy and Security Statement](#) | [Disclaimers](#)

Occupational Safety & Health Administration
200 Constitution Avenue, NW
Washington, DC 20210



ANTHRAX e-Tool Protecting the Worksite against terrorism

- [Home](#)
- [What is anthrax?](#)
- [Who is at risk?](#)
- [How do I prepare?](#)
- [What if I have a threat?](#)
- [How do I clean up my worksite?](#)

Additional Anthrax Response Information

The links associated with anthrax response are categorized as follows:



- Ⓞ [Comprehensive Anthrax Response References](#)
- Ⓞ [Mail Handling](#)
- Ⓞ [Health and Safety Plans / Medical Surveillance Programs](#)
- Ⓞ [Training and Personal Protective Equipment \(Code of Federal Regulations\)](#)
- Ⓞ [Sampling, Packing, and Transporting *Bacillus anthracis*](#)
- Ⓞ [Decontamination](#)

Comprehensive Anthrax Response References

- Ⓞ National Response Team, [Technical Assistance for Anthrax Response, Interim-Final Draft](#), October 2002.

Mail Handling

- Ⓞ Centers for Disease Control and Prevention (CDC) Health Advisory, [Updated Information About How To Recognize and Handle a Suspicious Package or Envelope](#), Distributed via the Health Alert Network October 31, 2001, CDCHAN-00050-01-10-31-ADV-N.
- Ⓞ Federal Bureau of Investigation (FBI) Advisory Poster, [If you receive a suspicious letter or package, what should you do?](#), General Information Bulletin 2000-3.
- Ⓞ United States Postal Service, [Mail Center Security Website](#).
- Ⓞ Public Health Foundation, Bioterrorism/Emergency Preparedness, [Video: Protecting Your Health for People Who Process, Sort, and Deliver the Mail](#)

Health and Safety Plans / Medical Surveillance Programs

- Ⓞ OSHA's *electronic health and safety plan template*
- Ⓞ Centers for Disease Control and Prevention (CDC), Morbidity and Mortality Weekly Report (MMWR), [Notice to Readers: Occupational Health Guidelines for Remediation Workers at *Bacillus anthracis*-Contaminated Sites --- United States, 2001--2002 \[September 6, 2002 / 51\(35\);786-789\]](#)
- Ⓞ American Conference of Governmental Industrial Hygienists (ACGIH), *2002 TLVs[®] and BEIs[®]* (extreme temperature reference), Publication #0102.

Training and Personal Protective Equipment (Code of Federal Regulations)

- ④ [29 CFR 1910.120, Hazardous Waste Operations and Emergency Response \(HAZWOPER\)](#)
- ④ [29 CFR 1910.38\(a\), Employee Emergency Plans and Fire Prevention Plans](#)
- ④ [29 CFR 1910.1200, Hazard communication](#)
- ④ [29 CFR 1910 Subpart I \(Sections 132 to 139\), Personal Protective Equipment](#)
- ④ 49 CFR 172 Subpart H, *Transportation Training*; consult the [Code of Federal Regulations website](#) for the most current version

Sampling, Packing, and Transporting *Bacillus anthracis*

- ④ Centers for Disease Control and Prevention, Public Health Emergency Preparedness and Response, [The Laboratory Response Network for Bioterrorism \(LRN\)](#) [a slideshow that provides a general overview of the LRN]
- ④ Centers for Disease Control and Prevention, Public Health Emergency Preparedness and Response, [Approved Tests for the Detection of *Bacillus anthracis* in the Laboratory Response Network \(LRN\)](#)
- ④ Centers for Disease Control and Prevention, Public Health Emergency Preparedness and Response, [Packing Critical Biological Agents](#) (protocols for packing, shipping, and transport of biological agents/diseases)
- ④ Centers for Disease Control and Prevention, Office of Health and Safety, Biosafety Branch, [Interstate Shipment of Etiologic Agents \(42 CFR Part 72\)](#)

Decontamination

- ④ The Environmental Protection Agency's (EPA's) Technology Innovation Office (TIO) has developed an information clearinghouse (www.EPATechBIT.org) as a centralized location to collect and disseminate information about decontamination technologies and also for technology vendors to provide information.

| [Home](#) | [What is Anthrax?](#) | [Who is at risk?](#) | [How do I prepare?](#) |
| [What if I have an anthrax threat?](#) | [How do I clean up my worksite?](#) |
| [Viewing/Printing Instructions](#) | [Credits](#) |

 [Back to Top](#)

www.osha.gov

www.dol.gov

[Contact Us](#) | [Freedom of Information Act](#) | [Information Quality](#) | [Customer Survey](#)
[Privacy and Security Statement](#) | [Disclaimers](#)

Occupational Safety & Health Administration
200 Constitution Avenue, NW
Washington, DC 20210